

CS 310: Algorithms


# Lecture 12

**Instructor:** Naveed Anwar Bhatti



#### Question 1:


Design an algorithm (pseudo or just steps) to check if a directed graph is strongly connected. Your algorithm **should only use Breadth-First Search (BFS)** and **should not calculate the reverse** (or transpose) of the graph G. After describing your algorithm, state its time complexity.





#### Question 1:


Design an algorithm (pseudo or just steps) to check if a directed graph is strongly connected. Your algorithm **should only use Breadth-First Search (BFS)** and **should not calculate the reverse** (or transpose) of the graph G. After describing your algorithm, state its time complexity.





#### Question 2:


Find topological order.






#### Question 2:


Find topological order.

































#### **Question 3:**


Proof through contradiction that Directed Acyclic Graph (DAG) cannot be a strongly connected graph.

- Assume for the sake of contradiction that there exists a DAG which is also strongly connected.
- Let's take any two vertices, **u** and **v**. Since the graph is assumed to be strongly connected, there exists a path from **u** to **v** and also from **v** to **u**.
- Since there exists a path from  $\mathbf{u}$  to  $\mathbf{v}$  and from  $\mathbf{v}$  to  $\mathbf{u}$ , it implies that we can start at vertex  $\mathbf{u}$ , travel to vertex  $\mathbf{v}$ , and then come back to vertex  $\mathbf{u}$ , forming a directed cycle.
- This contradicts our initial assumption that the graph is a DAG (as DAGs don't have directed cycles).
- Therefore, our initial assumption is wrong.



Midterm Exam on Monday, October 23<sup>rd</sup>





Chapter 5: **Divide and Conquer** 

Lower Bound on Sorting Algorithms

# RECORITHMS RECORDED TO THE PROPERTY OF THE PR

O, \O, and \O





Ali – (LUMS)

You, implement a sorting algorithm with worst-case runtime **O(n log log n)** by next week.




Okay Boss, I will try to do that ~






Thankyou,
You are FIRED






Ali try it for one month... can't do it

Boss, I can't do it







You, implement a sorting algorithm with worst-case runtime O(n log log n) by next week.



Jaffer – (LUMS)

No, Boss. O(n log log n) is below the lower bound on sorting algorithm complexity, I can't do it, nobody can do it!





## Lower bounds on sorting algorithms

#### **Know your limit**

we always try to make algorithms faster, but if there is a limit that you cannot exceed, you want to know

Is **O(n log n)** the best we can do?


Actually, yes, because the lower bound on sorting algorithms is  $\Omega(n \log n)$ , i.e., a sorting algorithm needs at least cn log n time to finish in worst-case.



## Lower bounds on sorting algorithms


# actually, more precisely ...

The lower bound **n** log **n** applies to only all **comparison based** sorting algorithms





# Comparison tree (for 3 distinct keys a, b, and c)



Slide credit: Kevin Wayne. Copyright © 2005 Pearson-Addise Wesley. All rights reserved.



## Sorting lower bound

Theorem. Any deterministic compare-based sorting algorithm must make  $\Omega(n \log n)$  compares in the worst-case.

#### Pf. [information theoretic]

- Assume array consists of n distinct values  $a_1$  through  $a_n$ .
- Worst-case number of compares = height h of pruned comparison tree.
- Binary tree of height h has  $\leq 2^h$  leaves.
- n! different orderings  $\Rightarrow n!$  reachable leaves.


$$2^{h} \geq \# \text{ reachable leaves} = n !$$

$$\Rightarrow h \geq \log_{2}(n!)$$

$$\geq n \log_{2} n - \ln(e)$$
Stirling's formula

Slide credit: Kevin Wayne. Copyright © 2005 Pearson-Addison Wesley. All rights reserved.





Chapter 5: **Divide and Conquer** 

- Use of Master Theorem
- Proof of Master Theorem (Next Lecture)

#### What is Master Theorem?

#### **Theorem**

If 
$$T(n) = aT\left(\frac{n}{b}\right) + f(n)$$
 (for constants **a**>0, **b**>1)

Then let T(n) has the following asymptotic bounds:

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$

• Using Master Theorem, find the asymptotic bounds of:

$$T(n) = 9T\left(\frac{n}{3}\right) + n$$

$$T(n) = aT(n/b) + f(n)$$

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$

• Using Master Theorem, find the asymptotic bounds of:

$$T(n) = T\left(\frac{2n}{3}\right) + 1$$

$$T(n) = aT(n/b) + f(n)$$

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \log n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$

• Using Master Theorem, find the asymptotic bounds of:

$$T(n) = 3T\left(\frac{n}{4}\right) + nlgn$$

$$T(n) = aT(n/b) + f(n)$$

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$

• Using Master Theorem, find the asymptotic bounds of:

$$T(n) = 2T\left(\frac{n}{2}\right) + \Theta(n)$$

$$T(n) = aT(n/b) + f(n)$$

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \log n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$

• Using Master Theorem, find the asymptotic bounds of:


$$T(n) = 8T\left(\frac{n}{2}\right) + \Theta(n^2)$$

$$T(n) = aT(n/b) + f(n)$$

- 1. If  $f(n) = O(n^{\log_b a \epsilon})$  for some constant  $\epsilon > 0$ , then  $T(n) = \Theta(n^{\log_b a})$
- 2. If  $f(n) = \Theta(n^{\log_b a})$ , then  $T(n) = \Theta(n^{\log_b a} \lg n)$
- 3. If  $f(n) = \Omega(n^{\log_b a + \epsilon})$  for some constant  $\epsilon > 0$ , and if  $af(n/b) \le cf(n)$ , for some constant c < 1 and all sufficiently large n, then  $T(n) = \Theta(f(n))$



# Thanks a lot



If you are taking a Nap, wake up.....Lecture Over